arXiv:1703.07102v1 [math.PR] 21 Mar 2017

AN EXPONENTIAL LIMIT SHAPE OF RANDOM q-PROPORTION BULGARIAN SOLITAIRE

Kimmo Eriksson
Mälardalen University, School of Education, Culture and Communication,
Box 883, SE-72123 Västerås, Sweden
kimmo.eriksson@mdh.se

Markus Jonsson[1]
Mälardalen University, School of Education, Culture and Communication,
Box 883, SE-72123 Västerås, Sweden
markus.jonsson@mdh.se

Jonas Sjöstrand
Royal Institute of Technology, Department of Mathematics,
SE-10044 Stockholm, Sweden
jonass@kth.se

March 22, 2017

Abstract

We introduce *p_n-random q_n-proportion Bulgarian solitaire* ($0 < p_n, q_n \leq 1$), played on n cards distributed in piles. In each pile, a number of cards equal to the proportion q_n of the pile size rounded upward to the nearest integer are candidates to be picked. Each candidate card is picked with probability p_n, independently of other candidate cards. This generalizes Popov's random Bulgarian solitaire, in which there is a single candidate card in each pile. Popov showed that a triangular limit shape is obtained for a fixed p as n tends to infinity. Here we let both p_n and q_n vary with n. We show that under the conditions $q_n^2 p_n n/\log n \to \infty$ and $p_n q_n \to 0$ as $n \to \infty$, the p_n-random q_n-proportion Bulgarian solitaire has an exponential limit shape.

1. Introduction

The game of Bulgarian solitaire has received a great deal of attention, see reviews by Hopkins [10] and Drensky [2]. The Bulgarian solitaire is played with a deck of n identical cards divided arbitrarily into a number of piles. A move consists of picking

[1]Corresponding author

a card from each pile and letting these cards form a new pile. If piles are sorted in order of decreasing size, every position in the solitaire is equivalent to a Young diagram of an integer partition of n.

Popov [15] considered a random version of Bulgarian solitaire defined by a probability $p \in (0, 1]$, such that one card from each pile is picked with probability p, independently of the other piles. We will refer to this stochastic process on configurations as *p-random* Bulgarian solitaire. The probabilities of configurations converge to a stationary distribution. Popov showed that as n grows to infinity and configuration diagrams are downscaled by $\sqrt{n}$ in both dimensions, the stationary probability of the set of configurations that deviate from a triangle with slope p by more than $\varepsilon > 0$ tends to zero. In this sense, random configurations has a limit shape.

The objective of the present paper is to study such limit shapes in a generalization of random Bulgarian solitaire.

1.1. q_n-proportion Bulgarian solitaire

Olson [13] introduced a generalization of Bulgarian solitaire in which the number of cards that are picked from a pile of size h is given by some non-negative valued function $\sigma(h)$. Eriksson, Jonsson and Sjöstrand [3] recently studied the special case when σ is well-behaved in the sense that $\sigma(1) = 1$ and both $\sigma(h)$ and $h - \sigma(h)$ are non-decreasing functions of h. In particular, they studied a special case that they called *q_n-proportion Bulgarian solitaire*, defined by the rule $\sigma(h) = \lceil q_n h \rceil$. This means that from each pile we pick a number of cards given by the proportion q_n of the pile size rounded upward to the nearest integer. To illustrate the effect of the parameter q_n, set it to 0.3 and consider the configuration $(6, 2, 2, 1)$. From the first pile we pick $\lceil 0.3 \times 6 \rceil = 2$ cards; similar calculations give that 1 card is picked from each of the other three piles. Note that for $q_n \leq 1/n$ exactly one card is always picked from each pile, retrieving the ordinary Bulgarian solitaire.

As n tends to infinity, Eriksson, Jonsson and Sjöstrand [3] determined limit shapes of stable configurations of q_n-proportion Bulgarian solitaire: In case $q_n^2 n \to 0$, the limit shape is triangular, which generalizes the limit shape result for the ordinary Bulgarian solitaire. For other asymptotic behavior of q_n, other limit shapes were obtained. Specifically, in case $q_n^2 n \to \infty$, the limit shape is exponential. The intermediate case $q_n^2 n \to C > 0$ produces a family of limit shapes interpolating between the triangular and the exponential shape.

1.1.1. p_n-random q_n-proportion Bulgarian solitaire

We shall examine a p_n-random version of q_n-proportion Bulgarian solitaire, in which the proportion q_n (rounded upward) of cards in a pile are only *candidates* to be picked, each of which is picked only with probability p_n, independently of all other

candidate cards. This process will be denoted by $\mathscr{B}(n, p_n, q_n)$. Note that in the special case of a fixed p and for $q_n \leq 1/n$, this process is equivalent to Popov's p-random Bulgarian solitaire.

Our focus will be on establishing a regime in which p_n-random q_n-proportion Bulgarian solitaire has an exponential limit shape.

2. The concept of limit shapes

In this section we give the precise definitions of the limit shapes we consider. Let $\mathcal{P}(n)$ be the set of integer partitions of n. For any partition $\lambda \in \mathcal{P}(n)$ with $N = N(\lambda)$ positive parts $\lambda_1 \geq \lambda_2 \geq \ldots \geq \lambda_N > 0$, define $\lambda_i = 0$ for $i > N(\lambda)$, and the *diagram* of λ as the Young diagram oriented such that the parts of λ are represented by left and bottom aligned columns, weakly decreasing in height from left to right. For example, is the diagram of the partition $(2, 1, 1)$. We define the *diagram-boundary function* of λ as the nonnegative, weakly decreasing and piecewise constant function $\partial\lambda : \mathbb{R}_{\geq 0} \to \mathbb{R}_{\geq 0}$ describing the boundary of λ, given by

$$\partial\lambda(x) = \lambda_{\lfloor x \rfloor + 1}.$$

Following [4] and [16], the diagram is downscaled using some *scaling factor* $a_n > 0$ such that all row lengths are multiplied by $1/a_n$ and all column heights are multiplied by a_n/n, yielding a constant area of 1. Following [3], we shall consistently make the choice $a_n = n/\lambda_1$, such that the height of the diagram is scaled to 1.

Thus, given a partition λ, define the *rescaled diagram-boundary function* of λ as the nonnegative, real-valued, weakly decreasing and piecewise constant function $\partial\lambda : \mathbb{R}_{\geq 0} \to \mathbb{R}_{\geq 0}$ given by

$$\partial\lambda(x) = \frac{1}{\lambda_1}\partial\lambda(xn/\lambda_1) = \frac{1}{\lambda_1}\lambda_{\lfloor xn/\lambda_1 \rfloor + 1}. \tag{1}$$

The p_n-random q_n-proportion Bulgarian solitaire $\mathscr{B}(n, p_n, q_n)$ (with $p_n, q_n \in (0, 1]$) can be regarded as a Markov chain on the finite state-space $\mathcal{P}(n)$. Let us denote the sequence of visited states by $(\lambda^{(0)}, \lambda^{(1)}, \ldots)$. In the truly random case of $p_n < 1$, it is straightforward to verify that this Markov chain is aperiodic and irreducible. It is well-known that an aperiodic and irreducible Markov chain on a finite state-space has a unique stationary distribution π and that starting from any initial state the distribution of the ith state $\lambda^{(i)}$ converges to π as i tends to infinity. We denote by π_{n,p_n,q_n} the stationary measure of the Markov chain $(\lambda^{(0)}, \lambda^{(1)}, \ldots)$ on $\mathcal{P}(n)$ given by $\mathscr{B}(n, p_n, q_n)$ for $p_n < 1$.[2] When we refer to a limit shape of the

[2]Readers acquainted with the limit shape literature may wonder whether the stationary measure has the property of being *multiplicative*, in the sense of interpretable as the product measure on the space of integer sequences restricted to a certain affine subspace [7]. The multiplicative property is useful in limit shape problems and related problems [1, 5, 8, 14, 16]. However, such techniques will not be used here as π_{n,p_n,q_n} is unlikely to be multiplicative in general.

process $\mathscr{B}(n, p_n, q_n)$ for $p_n < 1$ as n grows to infinity, we shall mean the limit shape of the stationary measure π_{n,p_n,q_n}. The intuitive sense of this concept is that *when the solitaire is played on a sufficiently large number of cards for sufficiently long the configuration will almost surely be very close to the limit shape after suitable downscaling*. Following Vershik [16], a sequence $\{\pi_n\}$ of probability distributions on $\mathcal{P}(n)$ is said to have a limit shape ϕ if the downscaled diagrams approach ϕ in probability as n grows to infinity. The exact condition for convergence can vary. Consistent with Yakubovich [17] and Eriksson and Sjöstrand [4], we shall use the definition that

$$\lim_{n\to\infty} \pi_n \left\{\lambda \in \mathcal{P}(n) : |\partial^{a_n}\lambda(x) - \phi(x)| < \varepsilon\right\} = 1 \tag{2}$$

for all $x > 0$ and all $\varepsilon > 0$.[3]

3. The approach of ordering piles by time of creation

It will sometimes be useful to explicitly order piles by time of creation rather than by size. Here we develop this approach.

When parts are not sorted by size, a configuration is not represented by an integer partition but by a *weak integer composition*: an infinite sequence $\alpha = (\alpha_1, \alpha_2, \dots)$, not necessarily decreasing, of nonnegative integers adding up to n. Let $\mathcal{W}(n)$ denote the set of weak compositions of n. We define the *diagram*, the *diagram-boundary function* $\partial\alpha$, and the *rescaled diagram-boundary function* $\partial\alpha$ of a weak composition α in exact analogy to the way we defined them for partitions in Section 2. For example, the diagram of $\alpha = (3, 0, 2, 4, 1, 0, 0, \dots)$ and the corresponding function graph $y = \partial\alpha(x)$ are shown in Figure 1. Also, for a weak composition $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_N, 0, 0, \dots)$ we define the number of parts $N = N(\alpha)$ disregarding the trailing zeros.

3.1. Connecting the limit shapes of compositions and partitions

We shall now connect compositions with partitions. For any $\alpha \in \mathcal{W}(n)$, define the operator ord as the ordering operator that arranges the parts of α in descending order, thus yielding a partition. We shall now prove that such sorting of the piles by size respects the convergence to a limit shape. The proof uses some basic theory

[3] Vershik [16] and Erlihson and Granovsky [6] used a stronger condition for convergence toward a limit shape, namely that

$$\lim_{n\to\infty} \pi_n \left\{\lambda \in \mathcal{P}(n) : \sup_{x\in[a,b]} |\partial^{a_n}\lambda(x) - \phi(x)| < \varepsilon\right\} = 1$$

should hold for any compact interval $[a, b]$, and any $\varepsilon > 0$.



Figure 1: The composition $\alpha = (3, 0, 2, 4, 1, 0, 0, \dots) \in \mathcal{W}(10)$.

of symmetric-decreasing rearrangements, see for example [9, Ch. 10] or [11, Ch. 3]. For any measurable function $f\colon \mathbb{R} \to \mathbb{R}_{\geq 0}$ such that $\lim_{x\to\pm\infty} f(x) = 0$, there is a unique function $f^*\colon \mathbb{R} \to \mathbb{R}_{\geq 0}$, called the *symmetric-decreasing rearrangement* of f, with the following properties:

- f^* is symmetric, that is, $f^*(-x) = f^*(x)$ for all x,
- f^* is weakly decreasing on the interval $[0, \infty)$,
- f^* and f are equimeasurable, that is,

$$\mathcal{L}(\{x :\ f(x) > t\}) = \mathcal{L}(\{x :\ f^*(x) > t\})$$

  for all $t > 0$, where $\mathcal{L}$ denotes the Lebesgue measure,

- f^* is lower semi-continuous.

In particular, if f is a symmetric function that is weakly decreasing and right-continuous on $[0, \infty)$ and tends to 0 at infinity, then $f^* = f$.

Lemma 1. *Let $\alpha \in \mathcal{W}(n)$ be a weak composition of n and let $f : \mathbb{R}_{\geq 0} \to \mathbb{R}_{\geq 0}$ be a right-continuous and weakly decreasing function such that $f(x) \to 0$ as $x \to \infty$. The downscaled diagram-boundary functions before and after sorting of the weak composition satisfy the inequality*

$$\|\partial \mathsf{ord}\, \alpha - f\|_\infty \leq \|\partial\alpha - f\|_\infty.$$

where $\|\cdot\|_\infty$ denotes the max-norm $\|f\|_\infty = \sup\big\{|f(x)| : x \geq 0\big\}$.

Proof. The intuition of the lemma should be obvious from Figure 2. To be able to use the standard machinery of symmetric rearrangements, we consider the functions

f, $\partial\alpha$, and $\partial\mathsf{ord}\,\alpha$ as being defined on the entire real axis by letting $f(x) = f(|x|)$ and analogously for $\partial\alpha$, and $\partial\mathsf{ord}\,\alpha$.

Since $f(x) \to 0$ as $x \to \infty$, its symmetric-decreasing rearrangement f^* is defined and, since f is weakly decreasing and lower semi-continuous, we have $f^* = f$. Similarly, $\partial\mathsf{ord}\,\alpha(x) \to 0$ as $x \to \infty$ and is weakly decreasing, so $(\partial\mathsf{ord}\,\alpha)^* = \partial\mathsf{ord}\,\alpha$. Moreover, $(\partial\alpha)^* = \partial\mathsf{ord}\,\alpha$ must hold because the operator ord arranges the composition parts in descending order.

Now, since symmetric rearrangements decrease L^p-distances for any $1 \le p \le \infty$ (see for example [11], Section 3.4), we obtain

$$\|\partial\mathsf{ord}\,\alpha - f\|_\infty = \|(\partial\alpha)^* - f^*\|_\infty \le \|\partial\alpha - f\|_\infty.$$

□



Figure 2: An example of a composition α and a decreasing function f showing that if $\partial\alpha(x)$ is enclosed between $f(x) - \varepsilon$ and $f(x) + \varepsilon$, then so is $\partial\mathsf{ord}\,\alpha(x)$, an immediate consequence of Lemma 1.

Clearly, Lemma 1 holds true also when the max-norm is replaced by the weaker convergence condition used in our limit shape definition (2).

Lemma 2. *For any distribution π_n on $\mathcal{W}(n)$, define a corresponding distribution $\tilde{\rho}^{(n)}$ on $\mathcal{P}(n)$ by*

$$\tilde{\rho}^{(n)}(\lambda) = \sum_{\substack{\alpha \in \mathcal{W}(n) \\ \mathsf{ord}\,\alpha = \lambda}} \pi_n(\alpha). \tag{3}$$

If ϕ is a limit shape of π_n on $\mathcal{W}(n)$ then ϕ is also a limit shape of $\tilde{\rho}^{(n)}$ on $\mathcal{P}(n)$.

Proof. The assumption that ϕ is a limit shape of the distribution π_n on $\mathcal{W}(n)$ means that

$$\lim_{n\to\infty} \pi_n \{\alpha \in \mathcal{W}(n) : |\partial\alpha(x) - \phi(x)| < \varepsilon\} = 1.$$

for all $x > 0$. By virtue of Lemma 1 we can replace α with $\mathsf{ord}\,\alpha$ in this formula:

$$\lim_{n\to\infty} \pi_n \{\alpha \in \mathcal{W}(n) : |(\partial \mathsf{ord}\,\alpha)(x) - \phi(x)| < \varepsilon\} = 1. \tag{4}$$

The set $A := \{\alpha \in \mathcal{W}(n) : |(\partial \mathsf{ord}\,\alpha)(x) - \phi(x)| < \varepsilon \text{ for all } x > 0\}$ can be written as a disjoint union of equivalence classes with respect to sorting:

$$A = \bigcup_{\lambda\in L} \{\alpha \in \mathcal{W}(n) : \mathsf{ord}\,\alpha = \lambda\}$$

where $L = \{\lambda \in \mathcal{P}(n) : |\partial\lambda(x) - \phi(x)| < \varepsilon \text{ for all } x > 0\}$. The π_n-probability measure of A is

$$\begin{aligned}
\pi_n(A) &= \pi_n\left(\bigcup_{\lambda\in L} \{\alpha \in \mathcal{W}(n) : \mathsf{ord}\,\alpha = \lambda\}\right) \\
&= \sum_{\lambda\in L} \pi_n\{\alpha \in \mathcal{W}(n) : \mathsf{ord}\,\alpha = \lambda\} \\
&= \sum_{\lambda\in L} \sum_{\substack{\alpha\in\mathcal{W}(n) \\ \mathsf{ord}\,\alpha=\lambda}} \pi_n(\alpha) \\
&= \sum_{\lambda\in L} \tilde{\rho}^{(n)}(\lambda) \qquad \text{(by (3))} \\
&= \tilde{\rho}^{(n)}(L).
\end{aligned}$$

From (4) we have that $\lim_{n\to\infty} \pi_n(A) = 1$. Because $\pi_n(A) = \tilde{\rho}^{(n)}(L)$, we can conclude that also $\lim_{n\to\infty} \tilde{\rho}^{(n)}(L) = 1$, that is,

$$\lim_{n\to\infty} \tilde{\rho}^{(n)} \{\lambda \in \mathcal{P}(n) : |\partial\lambda(x) - \phi(x)| < \varepsilon \text{ for all } x > 0\} = 1.$$

This means that ϕ is a limit shape of the distribution π_n on $\mathcal{P}(n)$. □

4. Three regimes

Recall from Section 1.1 the q_n-proportion Bulgarian solitaire developed in [3], where the limit shape is triangular when $q_n^2 n \to 0$, exponential when $q_n^2 n \to \infty$ and an interpolation between the two when $q_n^2 n \to C > 0$.

The p_n-random q_n-proportion Bulgarian solitaire seems to share this property of three regimes of limit shapes. Specifically, in Section 7 we conjecture the limit shape to be triangular when $p_n q_n^2 n \to 0$, exponential when $p_n q_n^2 n \to \infty$ and an interpolation between the two (a piecewise linear function graph that depends on C) when $p_n q_n^2 n \to C > 0$.

The focus in this paper is the exponential regime of the p_n-random q_n-candidate Bulgarian solitaire, i.e. the case $p_n q_n^2 n \to \infty$ as $n \to \infty$. However, with the proof technique we employ we will prove the stronger statement that the limit shape holds even when the configurations are considered elements of $\mathcal{W}(n)$, i.e. even without sorting the piles of a configuration according to size to create a partition in $\mathcal{P}(n)$. We will instead require the stronger condition $p_n q_n^2 n/\log n \to \infty$ as $n \to \infty$. By virtue of Lemma 2, the limit shape will also hold for partitions.

5. The exponential limit shape

Here we investigate the limit shape of configurations in the p_n-random q_n-candidate Bulgarian solitaire $\mathscr{B}(n, p_n, q_n)$ in the regime

$$\frac{p_n q_n^2 n}{\log n} \to \infty \text{ as } n \to \infty. \tag{5}$$

Our main result, Theorem 1, says that, under the additional asymptotic property $p_n q_n \to 0$ as $n \to \infty$, the boundary function of the diagram, downscaled, will resemble the exponential shape e^{-x} asymptotically almost surely (a.a.s.), i.e. with a probability that tends to 1 as $n \to \infty$. See Figure 3. Throughout this section, "a.a.s." can be read as "with a probability that tends to 1 as $n \to \infty$". Also, the asymptotic notations o and O will always be with respect to $n \to \infty$.



Figure 3: The result of a computer simulation after 200 moves of p_n-random q_n-proportion Bulgarian solitaire in the case $q_n = 1$, with $n = 10^5$ cards and $p_n = 0.01$, starting from a triangular configuration. The jagged curve is the rescaled diagram-boundary function of the resulting configuration and the smooth curve is the limit shape $y = e^{-x}$.

We shall see that the condition $p_n q_n^2 n/\log n \to \infty$ implies that the rounding effect in computing the number of candidate cards is negligible. Thus, the number of candidate cards will tend to $q_n n$ as $n \to \infty$. This in turn means that the expected number of picked cards will eventually be close to $p_n q_n n$, thus $\lambda_1 \approx p_n q_n n$ is the size of the pile created in a move of the solitaire. Recall from Section 2 that the scaling factor we employ is $a_n = n/\lambda_1 = \frac{1}{p_n q_n}$. Thus, if $p_n q_n$ is bounded away from zero, then the scaling $\frac{1}{p_n q_n}$ is bounded and hence cannot transform the jumpy boundary diagrams into a smooth limit shape. Therefore, we also require

$$p_n q_n \to 0 \text{ as } n \to \infty. \tag{6}$$

On the other hand, if $p_n q_n$ tends to zero too fast, the pile sizes will be small and their random fluctuations will be large. For instance, the new pile after each move has a size drawn from the binomial distribution $\mathrm{Bin}(K, p_n)$, where $K \approx q_n n$ is the number of candidate cards, with relative standard deviation $\sim 1/\sqrt{p_n q_n n}$. The requirement (5) guarantees that $p_n q_n$ does not tend to zero too fast.

Theorem 1. *For each positive integer n, pick q_n and p_n with $0 < p_n, q_n \leq 1$ and a (possibly random) initial configuration $\lambda^{(0)} \in \mathcal{P}(n)$. Let $(\lambda^{(0)}, \lambda^{(1)}, \dots)$ be the Markov chain on $\mathcal{P}(n)$ defined by $\mathscr{B}(n, p_n, q_n)$, and denote its stationary measure by π_{n,p_n,q_n}. Suppose*

$$p_n q_n \to 0 \quad \textit{and} \quad \frac{p_n q_n^2 n}{\log n} \to \infty \quad \textit{as } n \to \infty.$$

Then π_{n,p_n,q_n} has the limit shape e^{-x} under the scaling $a_n = (p_n q_n)^{-1}$.

The proof of Theorem 1 heavily relies on the following version of Chernoff bounds. For a proof, see for example [12].

Chernoff Bound. *For $n \geq 1$ and $0 < p \leq 1$, let $X \sim \mathrm{Bin}(n, p)$ and set $\mu = E(X) = np$. Then, for any $0 < \gamma < \mu$,*

$$P(|X - \mu| \geq \gamma) \leq 2 \exp\left(-\frac{\gamma^2}{3\mu}\right). \tag{7}$$

The idea of the proof of Theorem 1 is the following.

We will use the approach developed in Section 3, i.e. card configurations in the solitaire will be represented by weak integer compositions and the piles are ordered with respect to creation time, i.e. if $\alpha \in \mathcal{W}(n)$ is the current configuration in the solitaire, then α_1 was the last formed pile, α_2 the pile that was formed two moves ago, etc. With this representation, some piles may be empty, so one may imagine each pile being placed in a bowl and the bowls are lined up in a row on the table. In each move of the solitaire, the new (possibly empty) pile is put in a new bowl to the left of all old bowls. As mentioned in Section 4, we shall prove Theorem 1 as a limit

shape result for diagram-boundary functions of *compositions*. Thus, throughout this section, each configuration of n cards will be represented by an element of $\mathcal{W}(n)$. Also, in the following we may abbreviate $p = p_n$ and $q = q_n$ unless the dependency on n is crucial.

Assume a configuration $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_N, 0, 0, \dots)$ of n cards with $N = N(\alpha)$ piles (so that $\sum_{i=1}^{N} \alpha_i = n$) in the solitaire $\mathscr{B}(n, p, q)$. The number of candidate cards in the next move is $\kappa := \sum_{i=1}^{N} \lceil q\alpha_i \rceil$. We denote the rounding effect in pile $1 \le i \le N$ by $R_i := \lceil q\alpha_i \rceil - q\alpha_i$ and the total rounding effect by $R := \kappa - qn = \sum_{i=1}^{N} R_i$.

Clearly, $R < N$ (since $R_i < 1$ for any i), i.e. the total rounding effect is bounded above by the number of piles. The first thing we will do is to make sure that after a sufficient number D of moves from the initial configuration $\alpha^{(0)}$, the number of piles $N(\alpha^{(D)})$ in the resulting configuration $\alpha^{(D)}$ is much smaller than qn a.a.s. (so that the number of candidate cards κ is approximately qn and thus the total rounding effect R is negligible). In Lemma 4 we show that it is possible to choose such a D, namely $D = c\frac{\log n}{pq}$ for any $c \ge 14$.

We also need to make sure that the number of piles stays $o(qn)$ for sufficiently many additional moves M, long enough to establish the convergence of the overall shape. Lemma 4 will also guarantee that $M = \lceil n^2/p \rceil$ suffices for this purpose.

Thus, in the following we shall use

$$D = \left\lceil 14\frac{\log n}{pq} \right\rceil \quad \text{and} \quad M = \left\lceil \frac{n^2}{p} \right\rceil. \tag{8}$$

If the number of piles stays $o(qn)$ during M moves so that the number of candidate cards stays approximately qn, the newly formed pile in each of these moves will have expected size pqn. Our proof technique involves studying the evolution of such a pile (which will follow an exponential decay in size). Therefore we need to additionally make sure that no old piles (which could potentially be much larger than pqn) remain after these M moves. Lemma 3 shows that, in fact, after M moves all piles in the starting configuration have disappeared a.a.s.

Lemma 3. *Let M be given by* (8)*. From any initial configuration $\alpha \in \mathcal{W}(n)$, after M moves in the solitaire $\mathscr{B}(n, p_n, q_n)$, all piles in α have been consumed a.a.s.*

Proof. Consider a pile of size n. The size of this pile after M moves is statistically dominated by $\max(n - X, 0)$ where $X \sim \mathrm{Bin}(M, p_n)$ whose expected value is $E(X) = Mp_n = \lceil n^2/p_n \rceil p_n > n$. Therefore, the probability that the pile remains after M moves is $P(X < n)$ with the bound

$$\begin{aligned} P(X < n) &\le P(|X - Mp_n| > |Mp_n - n|) \le 2\exp\left(-\frac{(Mp_n - n)^2}{3Mp_n}\right) \\ &\le 2\exp\left(-\frac{n^2}{3}(1 + o(1))\right), \end{aligned}$$

where we used the Chernoff bound (7). Thus, since any given pile in α has size $\leq n$, and the number of piles (of any size in any configuration) is $\leq n$, the probability that *all* piles in α have been consumed after M moves is at least

$$1 - 2n \exp\left(-\frac{n^2}{3}(1+o(1))\right) \to 1$$

which concludes the proof. □

Lemma 4. *Let n, p_n, q_n and an initial configuration $\alpha^{(0)}$ be given in the solitaire $\mathscr{B}(n, p_n, q_n)$. Then*

$$\frac{1}{q_n n} \max\left\{N(\alpha^{(D+1)}), \ldots, N(\alpha^{(D+M)})\right\} \to 0 \text{ in probability,}$$

where D and M are given by (8).

Proof. Let us abbreviate $p = p_n$ and $q = q_n$. We will first prove that all piles of size at most $q^{-1}\log n$ disappear with high probability after D moves, making sure that there are not many small piles in $\alpha^{(D)}$. Consider a pile of size at most $q^{-1}\log n$ in $\alpha^{(0)}$. Note that every nonempty pile decreases by at least 1 with probability at least p in each move. Therefore, after D moves the number of picked cards from this pile statistically dominates $X \sim \mathrm{Bin}(D, p)$ with expected value $Dp = 14q^{-1}\log n$. Using the Chernoff bound (7), the probability that this pile remains after D moves is at most

$$\begin{aligned} P_1 &:= P\left(X < \frac{\log n}{q}\right) \leq P\left(|X - Dp| > \left|Dp - \frac{\log n}{q}\right|\right) \\ &\leq 2\exp\left(-\frac{\left(Dp - \frac{\log n}{q}\right)^2}{3Dp}\right) = 2\exp\left(-\frac{(\frac{14\log n}{q} - \frac{\log n}{q})^2}{3\cdot 14\frac{\log n}{q}}\right) = 2n^{-\frac{13^2}{42}\frac{1}{q}} < 2n^{-4}. \end{aligned}$$

Since there can be at most n piles of size at most $q^{-1}\log n$, the probability that not *all* piles of size at most $q^{-1}\log n$ have disappeared after D moves is bounded by

$$P_2 := nP_1 = 2n^{-3}.$$

Let us now turn our attention to *the number of* piles after these D moves. By the above, all piles smaller than $q^{-1}\log n$ have disappeared with high probability. Clearly, the number of piles *larger than* $q^{-1}\log n$ can never be more than $\frac{n}{q^{-1}\log n} = \frac{qn}{\log n}$. Also, during the process of these D moves, at most D new piles have been formed. (Exactly D piles have been formed but some may have disappeared in the process.) Thus, for the total number of piles $N(\alpha^{(D)})$ in the configuration $\alpha^{(D)}$ after D moves, with probability at least $1 - 2n^{-3}$, we have

$$N(\alpha^{(D)}) \leq \frac{qn}{\log n} + D = \frac{qn}{\log n} + 14\frac{\log n}{pq} = qn\left(\frac{1}{\log n} + 14\frac{\log n}{pq^2 n}\right) = o(qn),$$

where we used the assumption (5) in the last step. It follows that, for any $\varepsilon > 0$,

$$\frac{1}{qn} \max\left\{N(\alpha^{(D+1)}), \ldots, N(\alpha^{(D+M)})\right\} < \varepsilon$$

with probability at least $1 - 2n^{-3}M \geq 1 - \frac{2}{pn} \to 1$ since $pn \to \infty$. (That $pn \to 0$ is also a consequence of the assumption (5).) □

Lemma 4 asserts that the number of piles remains to be $o(qn)$ during the M moves from $\alpha^{(D)}$ to $\alpha^{(D+M)}$, hence the number of candidate cards remains to be qn (a.a.s.) during the same moves. Therefore the number of picked cards (which equals the size of the newly formed pile), remains of *expected* size pqn. In Lemma 5 we prove that the *actual* number of picked cards in each of these M moves does not deviate (relatively) from pqn.

Lemma 5. *Let n, p_n, q_n and an initial configuration $\alpha^{(0)}$ be given in the solitaire $\mathscr{B}(n, p_n, q_n)$. Let D and M be given by* (8)*. Then*

$$\max_{k \in [D+1, D+M]} \frac{|\alpha_1^{(k)} - p_n q_n n|}{p_n q_n n} \to 0 \text{ in probability}$$

as $n \to \infty$.

Proof. Let us abbreviate $p = p_n$ and $q = q_n$. Let $\varepsilon > 0$ and let κ be the number of candidate cards in $\alpha^{(k-1)}$ for some $k = D+1, \ldots, D+M$. Recall that the total rounding effect in computing the number of candidate cards is bounded above by the number of piles. It therefore follows from Lemma 4 that $\kappa = nq(1 + o(1))$. The new pile size is $\alpha_1^{(k)} \sim \mathrm{Bin}(\kappa, p)$. Then, using the triangle inequality and the Chernoff bound (7) we have

$$\begin{aligned} P_3 := P(|\alpha_1^{(k)} - pqn| > \varepsilon pqn) &\leq P(|\alpha_1^{(k)} - \kappa p| > \varepsilon pqn - |\kappa p - pqn|) \\ &< 2\exp\left(-\frac{(\varepsilon pqn - |\kappa p - pqn|)^2}{3\kappa p}\right) \\ &= 2\exp\left(-\frac{\varepsilon^2}{3} pqn(1 + o(1))\right) \\ &= o(1/M) \end{aligned}$$

where the last equality is derived as follows. By (5), $\log n = o(pqn)$ and hence $\log n^a = o(pqn)$ for any $a \geq 1$. Since $pqn \to \infty$, this means that $\exp(-pqn)$ tends to zero faster than $\exp(-\log n^a)$, i.e., $\exp(-pqn) = o(1/n^a)$. Since $np \to \infty$, we therefore also have $\exp(-pqn) = o(p/n^a) = o(1/M)$. The next to the last equality follows from the fact that εpqn dominates over $|pqn - \kappa p|$ (since $|pqn - \kappa p| = |pqn - nq(1 + o(1))p| = pqn \cdot o(1)$).

Therefore, the probability that $|\alpha_1^{(k)} - pqn| > \varepsilon pqn$ for *any* k during the entire process of M moves is bounded by $MP_3 = M \cdot o(1/M) = o(1)$. □

While playing the solitaire, there is a possibility that at some point there will be too many piles, and thereby the number of candidate cards will be bigger than qn (and thus the size of the newly formed pile will be bigger than pqn). Lemmas 3 and 4 ensures that this never happens a.a.s. during the entire process of M moves from $\alpha^{(D)}$ to $\alpha^{(D+M)}$.

There is also a risk that, even if there are suitably many (qn) candidate cards, the number of *picked* cards among them will deviate from pqn due to random fluctuations (and thereby the size of the newly formed pile will deviate from pqn). Lemma 5 ensures that this never happens a.a.s. during the same period of M moves.

Therefore, after $m := D + M$ moves we have the following a.a.s.

- all current piles have been formed during the last M moves, and
- all current piles had size pqn when they were formed.

At this point, i.e. in the configuration $\Gamma := \alpha^{(m)}$, the leftmost pile (of size Γ_1) was formed one move ago, the second pile from the left (of size Γ_2) was formed two moves ago, and so on. We shall prove that the size Γ_k of the pile that was formed k moves ago for any $k = 1, 2, \ldots, m$ is $\Gamma_k = \Gamma_1(1-pq)^k = pqn(1-pq)^k$ a.a.s., i.e. the size decreases exponentially with k with decay factor $1 - pq$.

We will now consider the evolution of a given pile of size A_1 during $r \geq 1$ steps in the p-random q-proportion Bulgarian solitaire in the following way. We will need to keep track of each individual card in this pile. To this end, we label the cards $1, 2, \ldots, A_1$ starting from the top, and each card will keep their label throughout the process. Let $X_{i,k} \in \{0, 1\}$ where $i = 1, \ldots, A_1$ and $k = 1, \ldots, r$ be independent Bernoulli random variables with $P(X_{i,k} = 1) = p$.

Consider the following process. Let A_{k+1} be the number of cards after k moves. In each move $k = 1, 2, \ldots, r$, we remove the card with label i if $X_{i,k} = 1$ and this card belongs to the candidate cards, i.e., the $\lceil qA_k \rceil$ top-most remaining cards. We will call this process a *q-process*. This process describes the evolution of a pile of size A_1 in the p-random q-proportion Bulgarian solitaire.

Using the same Bernoulli variables, for any real number $0 \leq s \leq 1$, we define an *s-threshold process* in the following way. In each move $k = 1, 2 \ldots, r$, we remove the card with label i if $X_{i,k} = 1$ and $i \leq sA_1$. In this process, we let $A_{k+1}^{[s]}$ denote the number of remaining cards after k moves. When it is relevant to indicate the initial pile size, an s-threshold process is called an (s, A_1)-threshold process and the number of remaining cards after k moves is denoted by $A_{k+1}^{[s,A_1]}$.

In the proof of Theorem 1, we will use two different s-threshold processes (for two different values of s) to over- and underestimate the sizes of $r + 1$ consecutive piles in Γ (corresponding to the r steps in an s-threshold process). Both these processes will have the same desired limit shape and thus the limit shape of our solitaire will follow by the squeeze theorem. We first need a combinatorial lemma giving sufficient conditions for overestimation and for underestimation.

Lemma 6. *(i) If* $\lceil sA_1 \rceil \le \lceil qA_1 \rceil$*, then* $A_k^{[s]} \ge A_k$ *for* $k = 1, \dots, r+1$.
(ii) If $(1-q)A_{r+1}^{[s]} \ge A_1 - \lceil sA_1 \rceil$*, then* $A_k^{[s]} \le A_k$ *for* $k = 1, \dots, r+1$.

Proof. (i) A card that is removed at some step ℓ during the s-threshold process must have label $i \le \lceil sA_1 \rceil$, so in the q-process it belongs to the $\lceil qA_1 \rceil$ candidate cards in the initial pile and hence it belongs to the candidate cards also at step ℓ and will be removed. Thus, every card removed in the s-threshold process is removed in the q-process too, and it follows that $A_k^{[s]} \ge A_k$ for $k = 1, \dots, r+1$.

(ii) We show by induction over r that, after r steps, the remaining cards in the s-threshold process is a subset of the remaining cards in the q-process. Suppose $(1-q)A_{r+1}^{[s]} \ge A_1 - \lceil sA_1 \rceil$. Since $A_r^{[s]} \ge A_{r+1}^{[s]}$ we have $(1-q)A_r^{[s]} \ge A_1 - \lceil sA_1 \rceil$ which by the induction hypothesis implies that $A_k^{[s]} \le A_k$ for $1 \le k \le r$. It follows that $(1-q)A_r \ge A_1 - \lceil sA_1 \rceil$ which in turn implies that $A_r - (A_1 - \lceil sA_1 \rceil) \ge \lceil qA_r \rceil$. This latter inequality means that the $\lceil qA_r \rceil$ topmost cards before step r in the q-process all have labels larger than $A_1 - \lceil sA_1 \rceil$. Thus, if a card is removed in step r in the q-process it is also removed in step r or earlier in the s-threshold process. This concludes the induction step. The base step $r = 0$ is trivial. □

Recall that we are considering the configuration $\Gamma = \alpha^{(m)}$ after $m = M + D$ moves in the solitaire from the initial configuration $\alpha^{(0)}$. We will compare the sizes of $r+1$ consecutive piles in Γ to the $r+1$ pile sizes in an s-threshold process. In order to make the comparison for *all* piles in Γ, this will be done for $r+1$ consecutive piles (which we will call an *r-chunk*) at a time. In each r-chunk the initial pile size is the corresponding pile size in the solitaire. In other words, $\Gamma_1, \Gamma_2, \dots, \Gamma_{r+1}$ will be compared to the pile sizes in an (s, Γ_1)-threshold process (with initial pile size Γ_1); and $\Gamma_{r+2}, \Gamma_{r+3}, \dots, \Gamma_{2(r+1)}$ will be compared to the pile sizes in an (s, Γ_{r+2})-threshold process (with initial pile size Γ_{r+2}), and so on. Let us call the resulting union of s-threshold processes an (r, s)-*union* process. Thus, if we denote the pile sizes in this (r, s)-union process by $U_1, U_2, \dots$, we have

$$\begin{aligned} &U_1 = \Gamma_1 = A_1^{[s,\Gamma_1]}, && U_2 = A_2^{[s,\Gamma_1]}, && \dots, && U_{r+1} = A_{r+1}^{[s,\Gamma_1]}, \\ &U_{r+2} = \Gamma_{r+2} = A_1^{[s,\Gamma_{r+2}]}, && U_{r+3} = A_2^{[s,\Gamma_{r+2}]}, && \dots, && U_{2(r+1)} = A_{r+1}^{[s,\Gamma_{r+2}]}, \dots . \end{aligned}$$

We intend to use the (r, s)-union process to estimate the pile sizes in Γ. In an s-threshold process, starting with a pile of size A_1, the number of remaining cards B above the level $A_1(1-s)$ after r moves is binomially distributed: $B \sim \mathrm{Bin}(A_1 s, (1-p)^r)$. See Figure 4. Therefore we need to choose $r = r_n$ and $s = s_n$ in such a way that we have the following in each s-threshold process:

I The pile size A_{k+1} is close to $A_1(1-pq)^k$ a.a.s. for all $k = 1, \dots, r$, which we need to establish the wanted limit shape.



Figure 4: The r steps of an (s, A_1)-threshold process.

II At the same time s must be close enough to q to make the over- and underestimations tight enough.

To accomplish (I), clearly $r = r_n$ can at least not be chosen bigger than $1/p_n$, in fact we shall require $p_n r_n \to 0$ as $n \to \infty$, in order for the variance in the size of the last pile (after r_n moves) in an s-threshold process to be small with high probability. However, we shall see that $p_n r_n$ may not tend to zero too fast. We will require

$$(p_n r_n)^2 \frac{p_n q_n^2 n}{\log n} = \frac{p_n^3 q_n^2 n r_n^2}{\log n} \to \infty \text{ as } n \to \infty. \tag{9}$$

(Recall from (5) that $p_n q_n^2 n / \log n \to \infty$.) However, since r_n is a positive integer for any n, if $p_n \not\to 0$ we cannot have $p_n r_n \to 0$, but will see that $r_n = 1$ suffices in the case $p_n \not\to 0$. In other words, we will require

$$p_n(r_n - 1) \to 0 \text{ as } n \to \infty. \tag{10}$$

To accomplish (II) we shall see that $s = q$ will suffice for the overestimation and $s = q(1 + 2pr) = q(1 + o(1))$ for the underestimation.

One way of choosing r_n such that (9) and (10) are fulfilled is

$$r_n = \left\lceil \rho_n^{-1/3} p_n^{-1} \right\rceil \text{ where } \rho_n = \frac{p_n q_n^2 n}{1 + \log n}. \tag{11}$$

This choice fulfills (10) since $(r_n - 1)p_n < r_n p_n \leq \rho_n^{-1/3} \to 0$. That (9) is fulfilled is easily verified:

$$(p_n r_n)^2 \frac{p_n q_n^2 n}{\log n} > \rho_n^{-2/3} \frac{p_n q_n^2 n}{\log n} = \frac{(p_n q_n^2 n)^{1/3}}{(1 + \log n)^{-2/3} \log n} > \left(\frac{p_n q_n^2 n}{\log n} \right)^{1/3} \to \infty$$

as $n \to \infty$ by (5).

Our next lemma, Lemma 7, will bound the probability P' that an initial pile of size $I_n := O(pqn)$ will, after r_n moves in an s_n-threshold process, deviate from the expected size assuming exponential decay, when $s_n = q_n(1 + o(1))$.

Since the number of piles is $\approx (p_n q_n)^{-1}$, the number of r-chunks is $\approx (p_n q_n r_n)^{-1}$. When using Lemma 7 we need the bound P' to hold for *each* chunk during *all* M moves (where M is given by (8)), specifically $P'M/(p_n q_n r_n) \to 0$ as $n \to \infty$. The probability in Lemma 7 is therefore bounded by $o(p_n q_n r_n/M) = o(p_n^2 q_n r_n/n^2)$. This is also why the pile size deviation $\varepsilon n p_n q_n$ is scaled with the number of chunks, resulting in the deviation $(\varepsilon p_n q_n n)(p_n q_n r_n) = \varepsilon p_n^2 q_n^2 n r_n$.

Lemma 7. *Let $(p_n)_n$ and $(q_n)_n$ be real sequences such that $0 < p_n, q_n \le 1$ and $p_n q_n \to 0$ as $n \to \infty$. For each n, let also $B_n \sim \mathrm{Bin}(F_n s_n, (1-p_n)^{r_n})$ where $(F_n)_n$ and $(s_n)_n$ are real sequences such that*

$$F_n = O(p_n q_n n) \quad \textit{and} \quad s_n = q_n(1 + o(1)), \tag{12}$$

and $F_n s_n$ is an integer for any n. Let also $(r_n)_n$ be the sequence of positive integers in (11)*. Then, for all $\varepsilon > 0$ we have*

$$P\left(\left|B_n + F_n(1-s) - F_n(1-p_n q_n)^{r_n}\right| > \varepsilon p_n^2 q_n^2 n r_n\right) = o(p_n^2 q_n r_n/n^2).$$

Proof. Let us abbreviate $F = F_n$, $B = B_n$, $p = p_n$, $q = q_n$, $r = r_n$ and $s = s_n$. Thus, we want to prove that

$$P := P\left(\left|B + F(1-s) - F(1-pq)^r\right| > \varepsilon p^2 q^2 n r\right) = o(p^2 q r/n^2).$$

We first note that the expected value $E(B) = Fs(1-p)^r$. Using the triangle inequality $|B + F(1-s) - F(1-pq)^r| \le |B - E(B)| + |E - F(1-pq)^r + F(1-s)|$ we obtain

$$P \le P(|B - E(B)| > \varepsilon p^2 q^2 n r - |E(B) - F(1-pq)^r + F(1-s)|).$$

By the Chernoff bound (7) we get

$$P \le 2\exp\left(-\frac{(\varepsilon p^2 q^2 n r - |E(B) - F(1-pq)^r + F(1-s)|)^2}{3E(B)}\right). \tag{13}$$

For the indices n for which $r_n > 1$ we have $rp \le 2(r-1)p \to 0$ and hence

$$(1-p)^r = 1 - pr + o(pr) \quad \text{and} \quad (1-pq)^r = 1 - pqr + o(pqr). \tag{14}$$

For the indices n for which $r_n = 1$, the relations in (14) are trivially true.

This means

$$\begin{aligned}|E(B)-F(1-pq)^r+F(1-s)| &= |Fs(1-p)^r - F(1-pq)^r + F(1-s)| \\ &= |Fs(1-pr+o(pr)) - F(1-pqr+o(pqr)) + F(1-s)| \\ &= F\Big(pr(q-s) + s\cdot o(pr) + o(pqr)\Big) \\ &= o(Fpqr) = o(p^2q^2nr). \qquad \text{(by (12))}\end{aligned}$$

Thus the numerator in (13) can be written $[(\varepsilon+o(1))p^2q^2nr]^2$. By the assumptions in (12), the denominator in (13) can be written

$$3E(B) = 3Fs(1-p)^r = 3\cdot O(pqn)\cdot q(1+o(1))\cdot O(1) = O(pq^2n).$$

Putting these together, the bound (13) on P can be written

$$-\frac{1}{\log P} = O\left(\frac{O(pq^2n)}{[(\varepsilon+o(1))p^2q^2nr]^2}\right) = O\left(\frac{1}{p^3q^2nr^2}\right) = o\left(\frac{1}{\log n}\right)$$

where (9) was used in the last step. Since $pqnr\to\infty$ (also by (9)) and $pqr\to 0$ (by (6) and (10)), we have $\frac{1}{pqr} = o(n)$ and hence $\log\frac{1}{pqr} = o(\log n)$. Therefore

$$-\frac{1}{\log P} = o\left(\frac{1}{\log n + \log\frac{1}{pqr}}\right) = o\left(\frac{1}{\log\frac{n}{pqr}}\right).$$

From this follows

$$\log P = o\left(\log\frac{pqr}{n}\right) = o\left(\log\frac{p^2qr}{n^2}\right),$$

thus $P = o(p^2qr/n^2)$. □

Note that Lemma 7 concerns an s-threshold process, i.e. only r steps. In other words, it asserts that

$$P\left(\left|A_{r+1} - A_1(1-p_nq_n)^{r_n}\right| > \varepsilon p_n^2q_n^2nr_n\right) = o(p_n^2q_nr_n/n^2), \tag{15}$$

where $A_1 = O(p_nq_nn)$ is the first pile size in an r-chunk and $A_{r+1} = (1-s_n)A_1 + B_n$ the last (see Figure 4). However, the deviation and the probability were chosen in such a way that they can be added over all r-chunks. This is done in Lemma 8 which bounds the probability for deviation for the *entire* union process. Specifically, we will show that, for any $C>0$, the piles in Γ formed at most $\frac{C}{pq}$ moves ago, i.e. Γ_k for $k\le\frac{C}{pq}$, will follow an exponential decay a.a.s. The sizes of the piles formed more than $\frac{C}{pq}$ moves ago ($k>\frac{C}{pq}$) will be shown to be sufficiently small to be close enough to the tail in the exponential limit shape.

Lemma 8. *Let $U_1, U_2, \ldots$ be the pile sizes in an (r_n, s_n)-union process corresponding to $\mathscr{B}(n, p_n, q_n)$, where the initial pile size is $U_1 = O(p_n q_n n)$, and r_n is given by (11) and $s_n = q_n(1 + o(1))$. Let $M = \lceil n^2/p_n \rceil$. Then*

$$\forall C, \varepsilon > 0 : \forall k < \tfrac{C}{p_n q_n} : P(|U_{k+1} - U_1(1 - p_n q_n)^k| > \varepsilon p_n q_n n) = o(1/M) = o(p_n/n^2).$$

Proof. As in the proof of Lemma 7, for the simplicity of notation we do not indicate in p, q, r and s the dependence on n. Let $C, \varepsilon > 0$ and $\varepsilon' = \varepsilon/C$. By the triangle inequality,

$$\begin{aligned}
&|U_{k+r+1} - U_1(1-pq)^{k+r}| \\
&\qquad \le |U_{k+1} - U_1(1-pq)^k|(1-pq)^r + |U_{k+r+1} - U_{k+1}(1-pq)^r| \\
&\qquad \le |U_{k+1} - U_1(1-pq)^k| + |U_{k+r+1} - U_{k+1}(1-pq)^r|.
\end{aligned}$$

Lemma 7 is now applicable for the first pile in each r-chunk (since $U_1 \ge U_2 \ge \cdots$ and $U_1 = O(p_n q_n n)$), so by its formulation (15), $|U_{k+r+1} - U_{k+1}(1-pq)^r| < \varepsilon' p^2 q^2 rn$ with probability $1 - o(p^2 qr/n^2)$. Thus,

$$|U_{k+r+1} - U_1(1-pq)^{k+r}| < |U_{k+1} - U_1(1-pq)^k| + \varepsilon' p^2 q^2 rn \tag{16}$$

with probability $1 - o(p^2 qr/n^2)$. We now note that the first term in the right hand side has the same form as the left hand side, only shifted with r piles. Thus, by induction it follows that, for any positive integer d, we have

$$|U_{dr+1} - U_1(1-pq)^{dr}| < d\varepsilon' p^2 q^2 rn$$

with probability $1 - o(p^2 qr/n^2)$. Thus, adding the probabilities for deviation for $k = r, 2r, \ldots, \eta r$, where $\eta = \lfloor \frac{C}{pqr} \rfloor$ we get

$$\begin{aligned}
&P\big(\forall k \in \{r, 2r, \ldots, \eta r\} : |U_{k+1} - U_1(1-pq)^k| > \eta\varepsilon' p^2 q^2 rn \ge \varepsilon pqn\big) \\
&\qquad = \eta \cdot o(p^2 qr/n^2) = o(p/n^2).
\end{aligned} \tag{17}$$

We have thereby proved the claim in the lemma for $k = r, 2r, \ldots, \eta r$. If k is not a multiple of r, suppose $dr < k < (d+1)r$ for some positive integer d. Then, since $pqr \to 0$ as $n \to \infty$ (which follows from (5) and (11)), we have $(1-pq)^r = 1 - pqr + o(pqr)$ and hence

$$|U_1(1-pq)^{(d+1)r} - U_1(1-pq)^{dr}| = O(pqn)(1-pq)^{dr}|pqr + o(pqr)| < \varepsilon pqn.$$

The lemma then follows by (17) and the fact that $U_{dr} \le U_k \le U_{(d+1)r}$. □

6. Proof of Theorem 1

Below follows the proof of Theorem 1, stated in Section 5.

Proof. First, as in the previous section, let us consider $\mathscr{B}(n, p_n, q_n)$ as a process on $\mathcal{W}(n)$ rather than on $\mathcal{P}(n)$, and let $\alpha^{(0)} \in \mathcal{W}(n)$ be the weak composition representing the initial configuration of cards in the solitaire. Let also M and D be given by (8).

Let $(r_n)_n$ be the sequence of positive integers given by (11) and let $(s_n)_n$ be the sequence $s_n = q_n(1 + 2p_n r_n)$. By Lemma 3 applied on $\alpha^{(D)}$, all piles present in $\alpha^{(D)}$ have disappeared in $\Gamma := \alpha^{(D+M)}$ a.a.s. Let $\Gamma_k = \alpha_k^{(D+M)}$ for $1 \le k \le M$ be the number of cards in the pile that was formed k moves ago. By Lemma 5, each of these piles had size $O(np_nq_n)$ a.a.s. when they were formed. Let $F_n := O(np_nq_n)$ be a sequence such that $F_n s_n$ is an integer for each n. Let $0 < \varepsilon < 1$ and choose C_n such that $C_n > \frac{p_n q_n \log \varepsilon}{\log(1-p_n q_n)}$.

Let $\check{U}_1, \check{U}_2, \dots$ be the pile sizes in the (r_n, s_n)-union process with initial pile size Γ_1. Using the fact that $p_n q_n r_n \to 0$, it is a straightforward computation to show that $s_n = q_n(1 + 2p_n r_n)$ implies $(1 - q_n)((1 - p_n r_n)^{r_n} - \varepsilon p_n q_n r_n) > 1 - s_n$ and therefore also

$$(1 - q_n)\big(A(1 - p_n r_n)^{r_n} - \varepsilon A p_n q_n r_n\big) > (1 - s_n)A \quad \text{for any } A > 0.$$

By Lemma 7, the probability that $\check{U}_1(1 - p_n r_n)^{r_n} - \varepsilon \check{U}_1 p_n q_n r_n < \check{U}_{r_n+1}$ is $P_1 := 1 - o(p_n^2 q_n r_n / n^2)$. Thus, with probability P_1 we have $(1-q_n)\check{U}_{r_n+1} > (1-s_n)\check{U}_1 \ge \check{U}_1 - \lceil s\check{U}_1 \rceil$ so by Lemma 6(ii), the pile sizes $\check{U}_1, \check{U}_2, \dots, \check{U}_{r_n+1}$ in the first r-chunk of the (r_n, s_n)-process underestimate the pile sizes $\Gamma_1, \Gamma_2, \dots, \Gamma_{r_n+1}$ with probability P_1. In the next chunk, we have a new absolute threshold $s\Gamma_{r_n+2} = s\check{U}_{r_n+2}$. Since $\Gamma_{r_n+2} \le \Gamma_1$, we have $(1 - q_n)\check{U}_{2r_n+2} > (1 - s_n)\check{U}_{r_n+2}$ with probability at least P_1, making Lemma 6(ii) applicable also for the second chunk to conclude that $\check{U}_{r_n+2}, \dots, \check{U}_{2r_n+2}$ underestimate $\Gamma_{r_n+2}, \dots, \Gamma_{2r_n+2}$ with probability at least P_1. Continuing in the same manner for the first $C_n(p_n q_n r_n)^{-1}$ chunks, we conclude that the (r_n, s_n)-union process underestimates the solitaire with high probability:

$$P(\check{U}_k > \Gamma_k \text{ for all } k < \frac{C_n}{p_n q_n}) < (1 - P_1)C_n(p_n q_n r_n)^{-1} = o(p_n/n^2).$$

Let $\widehat{U}_1, \widehat{U}_2, \dots$ be the pile sizes in the (r_n, q_n)-union process with initial pile size Γ_1. By Lemma 6(i) (with $s_n = q_n$), the (r_n, q_n)-union process surely overestimates the solitaire in each chunk.

Taking the results for the (r_n, s_n)-union process and the (r_n, q_n)-union process together we have

$$P(\check{U}_k \le \Gamma_k \le \widehat{U}_k \text{ for all } k < \frac{C_n}{p_n q_n}) > 1 - o(p/n^2).$$

Now, applying Lemma 8 to both the pile sizes $\check{U}_k$ and to the pile sizes $\widehat{U}_k$ and using the squeeze theorem, we obtain

$$\forall \varepsilon > 0 : \forall k < \tfrac{C_n}{p_n q_n} : P(|\Gamma_k - \Gamma_1(1-p_nq_n)^k| > \varepsilon n p_n q_n) < o(p_n/n^2).$$

Thus, the probability that $|\Gamma_k - \Gamma_1(1-p_nq_n)^k| < \varepsilon n p_n q_n$ for the first $\frac{C_n}{p_n q_n}$ piles throughout *all* M moves from $\alpha^{(D)}$ to $\alpha^{(D+M)}$ is $1 - M \cdot o(1/M) = 1 - o(1)$.

For piles $k > \frac{C_n}{p_n q_n}$, the exponential decrease (with decay factor $1-pq$) in pile size will yield piles smaller than $np_nq_n(1-p_nq_n)^{\frac{C_n}{p_nq_n}} < \varepsilon n p_n q_n$ (by our choice of C_n). Thus, the pile sizes themselves are below $\varepsilon n p_n q_n$.

In summary; playing sufficiently many moves of $\mathscr{B}(n, p_n, q_n)$, the resulting composition diagram will a.a.s. be arbitrarily close to the boundary diagram of the composition α where $\alpha_k = np_nq_n(1-p_nq_n)^{k-1}$ for all $k = 1, 2, \dots$. The corresponding boundary function is $\partial\alpha(x) = np_nq_n(1-np_nq_n)^{\lfloor x \rfloor}$. The corresponding rescaled boundary function, with the given scaling factor $a_n = (p_nq_n)^{-1}$, is

$$\partial^{a_n}\alpha(x) = (1-p_nq_n)^{\frac{x}{p_nq_n}} \to e^{-x}$$

since $p_nq_n \to 0$ as $n \to \infty$.

Setting $m := D+M$, and letting π_n^m denote the probability distribution on $\mathcal{W}(n)$ for $\alpha^{(m)}$, we have

$$\lim_{n\to\infty} \pi_n^m \left\{\alpha \in \mathcal{W}(n) : |\partial^{a_n}\alpha(x) - e^{-x}| < \varepsilon\right\} = 1,$$

for all $\varepsilon > 0$ and all $x > 0$, in accordance with (2). By virtue of Lemma 2, the same limit shape holds when configurations in the solitaire $\mathscr{B}(n, p_n, q_n)$ are represented by partitions $\mathcal{P}(n)$.

Since π_{n,p_n,q_n} is the stationary distribution of the Markov chain $(\lambda^{(0)}, \lambda^{(1)}, \dots)$, if we start with a partition $\lambda^{(0)}$ sampled from π_{n,p_n,q_n} and play m moves, the resulting partition $\lambda^{(m)}$ will also be sampled from π_{n,p_n,q_n}. Thus, the theorem follows by choosing $\lambda^{(0)}$ as a stochastic partition sampled from the stationary distribution. □

7. Conjectures

Recall that Theorem 1 was proved with $\mathscr{B}(n, p_n, q_n)$ being considered a process on $\mathcal{W}(n)$, and by virtue of Lemma 2 it also holds in $\mathcal{P}(n)$. We imposed the condition $\frac{p_nq_n^2 n}{\log n} \to \infty$. Here we conjecture that the weaker condition $p_nq_n^2 n \to \infty$ suffices in order for Theorem 1 to hold in $\mathcal{P}(n)$.

Conjecture 1. *Theorem 1 holds also when the condition $np_nq_n^2/\log n \to \infty$ is replaced by the weaker condition $np_nq_n^2 \to \infty$.*

The reason for this conjecture can be understood by considering the example $q_n = 1$ and $p_n n = \log(\log n)$. For this example it is easy to prove that there is no limit shape when sorting is not performed. Since $q_n = 1$, the number of picked cards in each move and thus the expected size of a new pile is $\mathrm{Bin}(n, p_n)$ with expected value np_n and standard deviation $\sigma \approx \sqrt{np_n}$. A pile of size np_n will after $1/p_n$ moves have the expected size $np_n(1-p_n)^{1/p_n} \to e^{-1}np_n$ as $n \to \infty$.

Thus, the probability for a "visible" deviation (i.e. greater than $d = \sqrt{np_n}$ standard deviations) is $P(\text{deviation} \geq d\sigma) = e^{-np_n}$, so for $1/p_n$ piles, the probability for a visible deviation anywhere is $\frac{e^{-np_n}}{p_n} = \frac{n}{\log n \log(\log n)} \to \infty$ as $n \to \infty$, i.e. the expected number of such large deviations tends to infinity as n tends to infinity. This makes it impossible to achieve a convergence in probability towards a limit shape. However, from simulations we have reason to believe that the process converges towards a limit shape when sorting is performed.

Further, recall from Section 4 the other regimes $np_nq_n^2 \to 0$ and $np_nq_n^2 \to C$ from some constant $C > 0$. We conjecture that the limit shapes in the p_n-random q_n-proportion Bulgarian solitaire in these regimes are the same as in the deterministic q-proportion Bulgarian solitaire developed in [3].

Conjecture 2. *If $p_nq_n^2n \to 0$ as $n \to \infty$, the limit shape of the p_n-random q_n-proportion Bulgarian solitaire is triangular.*

Conjecture 3. *If $p_nq_n^2n \to C$ as $n \to \infty$ for some constant $C > 0$, the limit shape of the p_n-random q_n-proportion Bulgarian solitaire is a piecewise linear shape that depends on the value of C.*

References

[1] S. Corteel, B. Pittel, C.D. Savage, and H.S. Wilf, *On the multiplicity of parts in a random partition*, Random Structures Algorithms **14** (1999), no. 2, 185–197.

[2] V. Drensky, *The Bulgarian solitaire and the mathematics around it*, Math. and Education in Math., Proc. of the 44-th Spring Conf. of the Union of Bulgar. Mathematicians, SOK-Kamchia, April 2–6, 2015 (2015), 79–91.

[3] K. Eriksson, M. Jonsson, and J. Sjöstrand, *Limit shapes of stable configurations of a generalized Bulgarian solitaire*, (forthcoming).

[4] K. Eriksson and J. Sjöstrand, *Limiting shapes of birth-and-death processes on Young diagrams*, Adv. in Appl. Math. **48** (2012), no. 4, 575 – 602.

[5] M. Erlihson and B. Granovsky, *Reversible coagulation–fragmentation processes and random combinatorial structures: Asymptotics for the number of groups*, Random Structures Algorithms **25** (2004), no. 2, 227–245.

[6] ———, *Limit shapes of Gibbs distributions on the set of integer partitions: The expansive case*, Ann. Inst. H. Poincaré Probab. Stat. **44** (2008), 915–945.

[7] B. Fristedt, *The structure of random partitions of large integers*, Trans. Amer. Math. Soc. **337** (1993), no. 2, 703–735.

[8] W.M.Y. Goh and P. Hitczenko, *Random partitions with restricted part sizes*, Random Structures Algorithms **32** (2008), no. 4, 440–462.

[9] G.H. Hardy, J.E. Littlewood, and G. Pólya, *Inequalities*, Cambridge University Press, Cambridge, 1934.

[10] B. Hopkins, *30 years of Bulgarian solitaire*, College Math. J. **43** (2012), no. 2, 136–141.

[11] E.H. Lieb and M. Loss, *Analysis, volume 14 of graduate studies in mathematics*, vol. 4, American Mathematical Society, Providence, RI, 2001.

[12] M. Mitzenmacher and E. Upfal, *Probability and computing: Randomized algorithms and probabilistic analysis*, Cambridge University Press, 2005.

[13] J.S. Olson, *Variants of Bulgarian solitaire*, Integers **16** (2016), A8.

[14] B. Pittel, *On a likely shape of the random Ferrers diagram*, Adv. in Appl. Math. **18** (1997), no. 4, 432–488.

[15] S. Popov, *Random Bulgarian solitaire*, Random Structures Algorithms **27** (2005), no. 3, 310–330.

[16] A. Vershik, *Statistical mechanics of combinatorial partitions, and their limit shapes*, Funct. Anal. Appl. **30** (1996), no. 2, 90–105.

[17] Y. Yakubovich, *Ergodicity of multiplicative statistics*, J. Combin. Theory, Series A **119** (2012), no. 6, 1250–1279.